

DATA SHEET

ANTI HUMAN PLASMINOGEN ACTIVATOR INHIBITOR 2 [PAI-2] ANTIBODY, POLYCLONAL

Catalog no.:	A 7631.1 / A 7631.2
Immunogen:	purified human PAI-2
Reference no.:	LocusID 5055
Host:	Goat
Matrix:	Serum
Specificity:	Human PAI 2. There were no cross reactivities obtained with human tPA ; uPA; and PAI-1.
Contents:	20 µl / 100 µl (lyophilized) resuspend in 20 µl / 100 µl aqua bidest
Known applications:	ELISA (1:5000), Westernblot (1:100), Immunohistochemistry, paraffin sections (5-10 µg/ml), cryo sections This antibody has not been tested for use in all applications. This does not necessarily exclude its use for non-tested procedures. The stated dilutions are recommendations only. We suggest that the applicant titrates the antibody in his/her system using appropriate negative/positive controls.
Store at:	2-8 °C (lyophilized); - 20 °C (dissolved) Repeated thawing and freezing must be avoided
References:	<ol style="list-style-type: none">1. Reinartz J, Schaefer B, Bechtel MJ, Kramer MD. Plasminogen activator inhibitor type-2 (PAI-2) in human keratinocytes regulates pericellular urokinase-type plasminogen activator. <i>Exp Cell Res.</i> 1996. 223(1):91-101.2. Ferrier CM, van Geloof WL, de Witte HH, Kramer MD, Ruiten DJ, van Muijen GN. Epitopes of components of the plasminogen activation system are re exposed in formalin-fixed paraffin sections by different retrieval techniques. <i>J Histochem Cytochem.</i> 1998. 46(4):469-476.3. Bechtel MJ, Wysocki NS, Heidtmann A, Stark HJ, Fusenig N, Kramer MD, Schaefer BM. Plasminogen activator inhibitor type 2 is expressed in keratinocytes during re-epithelialization of epidermal defects. <i>Br J Dermatol.</i> 1998. 138(1):22-28

Only for research use

N:\WINWORD.DAT\DATENBLAETTER\ANTIKOERPER\ONCOLOG\YA 7631.DOC