



## Data Sheet



# HUMAN UROKINASE PLASMINOGEN ACTIVATOR-RECEPTOR [UPA-R]

## ANTIBODY, POLYCLONAL

<b>Catalog no.:</b>	A 7612.1 / A 7612.2
<b>Immunogen:</b>	Purified human Urokinase Plasminogen Activator-Receptor
<b>Synonyms:</b>	Urokinase plasminogen activator surface receptor, U-PAR, uPAR, Monocyte activation antigen Mo3, CD87
<b>Swiss-Prot No:</b>	Q03405
<b>Gene Information:</b>	Gene Name: PLAUR, MO3, UPAR GeneID: 5329
<b>Host:</b>	Goat
<b>Matrix:</b>	Serum
<b>Specificity:</b>	Human Urokinase Plasminogen Activator-Receptor There were no cross reactivities obtained with human tPA and human PAI-2
<b>Contents:</b>	20 µl / 100 µl (lyophilized) Resuspend in 20 µl / 100 µl aqua bidest.
<b>Known applications:</b>	ELISA, Immunohistochemistry (Frozen sections & paraffin sections) Recommended procedure for immunohistochemistry: Dilutions for frozen sections: 1:50 – 1: 100 Dilutions for paraffin sections: 1:50 Recommended procedure for paraffin sections: Pretreatment with 0.05% Pronase E (Sigma) for 20 minutes. <small>This antibody has not been tested for use in all applications. This does not necessarily exclude its use in non-tested procedures. The stated dilutions are recommendations only. End users should determine optimal dilutions in their system using appropriate negative/positive controls.</small>
<b>Store at:</b>	2-8 °C (lyophilized); - 20 °C (dissolved) Repeated thawing and freezing must be avoided
<b>Last updated on:</b>	07 April 2022



Antibodies



**For research use only**

**Publishing research using A 7612? Please let us know so that we can cite your publication as a reference.**



**Immundiagnostik AG**

Stubenwald-Allee 8a · 64625 Bensheim · Germany

Phone: +49 6251 70190-0 · Fax: +49 6251 70190-363 · dept.immuochemicals@immundiagnostik.com · www.immundiagnostik.com